

ORGANIC CHEMISTRY-III

UNIT 1 NOTES

OPTICAL ISOMERISM

- OPTICAL ACTIVITY
- ENANTIOMERISM
- DIASTEREOMERISM
- MESO COMPOUNDS
- ELEMENTS OF SYMMETRY
- CHIRAL & ACHIRAL MOLECULES
- DL SYSTEM OF NOMENCLATURE
- RS SYSTEM OF NOMENCLATURE
- REACTION OF CHIRAL MOLECULES
- RECEMIC MODIFICATION
- ASYMMETRIC SYNTHESIS



CONNECT WITH US ON :




IMPERFECT PHARMACY APP




@IMPERFECTPHARMACY

STEREISOMERISM


- The Branch of Chemistry dealing with structure of three dimensional molecules is known as Stereochemistry.
- The phenomenon of isomerism caused by different arrangement of atoms or groups in space is known as Stereoisomerism, and the compounds showing phenomenon is known as Stereoisomers.
- These molecules have same structural as well as molecular formulas, but their atomic arrangement is different.



trans - 1,2 - dichloroethane



cis - 1,2, dichloroethane




OPTICAL ISOMERISM

- Optical Isomerism is a type of stereoisomerism in which molecules have the same molecular and structural formula but differ in the way they interact with plane-polarised light.
- They are also known as Enantiomers.

OPTICAL ACTIVITY


- When a plane polarised light is passed through an Optically Active Substance then it rotates the light in Clockwise or Anticlockwise direction and this phenomenon is known as Optical Rotation .
- The compounds showing Optical Activity are known as Optically Active Compounds.



- Now, if the compound rotates the light in Clockwise direction, then it is said to be Dextrorotatory (+)
- If the compound rotates the light in Anticlockwise direction, then it is said to be Levorotatory (-)
- If the light passes undeviatedly, then compound is Optically Inactive.

ENANTIOMERISM


- Enantiomerism is nothing but Optical Isomerism
- Enantioisomerism is a form of Stereoisomerism in which molecules exist as non-superimposable mirror images of each other.
- These mirror image isomers are called Enantiomers.



- Now, If the compound showing enantiomerism rotates the plane polarised light in clockwise direction, then it is said to be Dextrorotatory (+).
- And if in anticlockwise direction, then it is said to be Levorotatory (-).

DIASTEREOISOMERISM

- Diastereoisomerism is a type of Stereoisomerism where isomers have same molecular formula and connectivity of atoms but are not mirror images of each other and not superimposable.
- Unlike, Enantiomers, diastereoisomers do not have identical physical or chemical properties and often exhibit different melting points, boiling points and reactivity.




DIFFERENCE BETWEEN ENANTIOMERS & DIASTEREOISOMERS

PARAMETER	ENANTIOMER	DIASTEREOMER
• Number of Chiral Centres	One	Two or More
• Mirror Images	Yes	No
• Superimposition	No	No
• Physical Properties	Same	Different
• Chemical Properties	Same	Different

MESO COMPOUND

- A Meso compound is a stereoisomer that contains two or more chiral centers but is optically inactive due to an internal plane of symmetry.
- A meso compound can be divided into two symmetrical halves, making one half a mirror image of the other.
- This symmetry causes the molecule to exhibit internal compensation, meaning that the optical rotation of chiral centres cancel each other out.




TARTARIC ACID (MESO FORM)

ELEMENTS OF SYMMETRY



- Elements of symmetry refers to the geometric transformations in which you can move or change the position or shape of an object so that it still look the same.
- If certain symmetry elements are present, the molecule cannot show Optical Isomerism.
- The key symmetry elements are :
 - ① Plane of Symmetry
 - ② Centre of Symmetry
 - ③ Axis of Symmetry

① PLANE OF SYMMETRY


- A Plane of symmetry is an imaginary flat line that divides a molecule or object into two mirror-images halves.
- If you fold the object along this plane, both sides would match perfectly.
- In Optical Isomerism, if a molecule has a plane of symmetry, it means the molecule is achiral.



FOR LECTURE VIDEOS VISIT OUR  CHANNEL




FOR MORE CONTENT : VISIT OUR  APP




② CENTRE OF SYMMETRY

- A Centre of Symmetry (also known as Point of Symmetry) is a point in a molecule or object where every part has an identical counterpart in the opposite direction at the same distance.
- If a molecule has centre of symmetry it cannot be chiral.



③ AXIS OF SYMMETRY

An Axis of symmetry is an imaginary line around which an object or molecule can be rotated by a certain angle and it still looks the same.



CHIRAL MOLECULES

- A molecule is Chiral if it cannot be superimposed on its mirror image - just like left and right hands.
- These molecules lack a plane of symmetry and usually have a Chiral Centre (a carbon atom bonded to 4 ~~carbon~~ different groups)
- These molecules are Optically Active

ACHIRAL MOLECULES


- A molecule is achiral if it can be superimposed on its mirror image.
- These molecules have a plane of symmetry or centre of symmetry making them Optically Inactive.

DL SYSTEM OF NOMENCLATURE


The DL System of Nomenclature is a way to designate the configuration of chiral molecules, particularly amino acids and sugars, based on their similarity to the reference molecule glyceraldehyde.

RULES FOR DL SYSTEM

- Draw the Fischer Projection for the given molecule.
- Arrange the molecules so that the most oxidised group is at the top.
- Now look at the last chiral (asymmetric) carbon in the chain.
- Now check the position of OH, NH₂ groups
- If the -OH group on the last chiral carbon is on the RIGHT, the molecule is in the D- configuration.
- If the -OH group is on the LEFT, the molecule is in the L- configuration.



D - Glucose



L- Glucose

EXAMPLE

- D- Glucose : The -OH on the last chiral carbon is on right.
- L- Glucose : The -OH on the last chiral carbon is on left.

KEY FEATURES

- It does not correspond to optical activity (+ or -)
- Some D compounds are dextrorotatory (+) & some are levorotatory (-).
- Example : D- glucose is (+) but D- fructose is (-)

IMPORTANT NOTE


- DL System basically a Relative Configuration.
- The D/L system is an important convention in Organic Chemistry, especially for naming sugars and amino acids, but it is now often replaced by R/S system in modern chemistry.

RS SYSTEM OF NOMENCLATURE

- The R/S system is a modern and systematic way of naming chiral molecules based on spatial arrangement of atoms around a chiral centre.
- It was developed by Cahn, Ingold and Prelog and is used to determine the absolute configuration of Chiral Molecules.
- Unlike the D/L system, which is specific to sugars & amino acids, the R/S system can be applied to all chiral compounds.


RULES FOR R/S SYSTEM

- Identify the Chiral Center.
- Assigned priorities based on atomic number.
- The higher the atomic number, the higher the priority.
- Example: $\text{Br} > \text{Cl} > \text{O} > \text{N} > \text{C} > \text{H}$.
- If the first atom in the two groups is the same, look at the next atom in the chain until a difference is found.
- Double bond or triple bond is treated as if the atom is duplicated or triplicated.
- The lowest priority group should be on vertical line.
- Draw a curved arrow from priority 1 $\rightarrow$ 2 $\rightarrow$ 3.
- If the arrow moves clockwise, the configuration is R (Rectus).
- If the arrow moves anticlockwise, the configuration is S (Sinister).



SPECIAL CASES

- If the lowest priority group lies on horizontal plane instead of vertical then whatever configuration comes, just reverse it.
- If rotation appears R, assign S
- If rotation appears S, assign R



Importance Of RS System


- It is a universal system that applies to all chiral molecules not just biomolecules.
- It provides an absolute configuration rather than a relative one.

REACTIONS OF CHIRAL MOLECULES

- Chiral molecules can undergo various chemical reactions, often leading to products with specific stereochemistry.
- Chiral Molecules generally give 3 types of chemical reactions.
 - ① Retention
 - ② Inversion
 - ③ Racemization



① RETENTION

Retention reaction refers to a reaction where the configuration (R or S) of chiral centre remains the same after the reaction. means stereochemistry of chiral molecule is retained.




② INVERSION

An Inversion reaction in a chiral molecules refers to the process where the configuration (R or S) of a chiral centre is reversed after the reaction.




② - 2-Bromobutane




③ - 2-Butanol

③ RECEMIZATION / RECEMIC MODIFICATION

- Racemization is a process in which an optically active compound converts into a racemic mixture, which contains equal amount of both enantiomers (R & S form).
- Since the two enantiomers rotate light in opposite directions, the net optical activity becomes zero.



① (R) - 2 - Bromobutane



② (R) - 2 - Butanol

③ (S) - 2 - Butanol

RESOLUTION OF RACEMIC MIXTURE

- The Resolution of racemic mixture refers to the process of separating a racemic (50:50) mixture of enantiomers into its individual optically active components.
- It involves following techniques:

① Mechanical Separation

- Mechanical separation is one of the earliest methods used to resolve racemic mixtures.
- This technique relies on physically separating the crystals of individual enantiomers based on different shapes.

Process

- The mixture is allowed to crystallize slowly under controlled conditions.
- The crystals are examined under microscope.
- The individual enantiomeric crystals are picked out using a fine needle or tweezers.

② CHEMICAL SEPARATION

- Chemical separation is one of the most commonly used methods to separate a racemic mixture.
- In this a chiral resolving agent is used to convert enantiomers into diastereomers.
- Diastereomers have different physical properties, allowing separation via crystallization or chromatography.
- The resolving agent is then removed to obtain pure enantiomers.

③ ENZYMATIC SEPARATION


- Enzymatic separation is a highly selective method for separating a racemic mixture into its individual enantiomers using enzymes.
- Enzymes are biological catalysts that often exhibit stereoselectivity, meaning they react preferentially with only one enantiomer while leaving the other unchanged.

④ CHROMATOGRAPHIC SEPARATION

- Chromatographic separation is a powerful method for separating enantiomers from a racemic mixture using Chiral Stationary Phases in HPLC, TLC or Gas Chromatography.
- One enantiomer binds more strongly to CSP, while the other elutes faster.

ASYMMETRIC SYNTHESIS

- Asymmetric Synthesis is defined as conversion of an achiral (symmetric) compound into a chiral (asymmetric) compound in such a way that one enantiomer predominates over the other.




TYPES OF ASYMMETRIC SYNTHESIS

They are of two types :

- ① Partial Asymmetric Synthesis
- ② Absolute Asymmetric Synthesis


① PARTIAL ASYMMETRIC SYNTHESIS

- Partial asymmetric synthesis produces a chiral product with an excess of one enantiomer, but not exclusively.
- A mixture of enantiomer is formed with one being more abundant.



② Absolute ASYMMETRIC SYNTHESIS

Absolute Asymmetric synthesis creates only 1 specific enantiomer directly from an achiral material.



THANK YOU

FOR CHOOSING IMPERFECT PHARMACY AS YOUR STUDY PARTNER



JOIN US ON :



IMPERFECT PHARMACY



IMPERFECT PHARMACY



@IMPERFECTPHARMA



IMPERFECT PHARMACY



@IMPERFECTPHARMACY



IMPERFECT PHARMACY



IMPERFECTPHARMACY@GMAIL.COM